

GALAXIES AT $Z > 4.5$

ESTHER M. HU

*Institute for Astronomy, University of Hawaii, 2680 Woodlawn Dr.,
Honolulu, HI 96822, USA*

RICHARD G. MCMAHON

Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, UK

EIICHI EGAMI

*Max-Planck-Institut für extraterrestrische Physik, Postfach 1603,
85740 Garching, GERMANY*

We present the results of ground-based and HST imaging studies targeted on $z > 4.5$ quasar fields. High-redshift galaxies identified in deep narrow-band Ly α images of the fields surrounding the quasars BR1202–0725 ($z = 4.694$) and BR2237–0607 ($z = 4.558$) have been confirmed with follow-up spectroscopy using the LRIS spectrograph on the Keck 10m telescope. These high-redshift galaxies are typically only moderately luminous objects and their sub- L^* magnitudes imply star formation rates of only a few solar masses per year. For BR1202–0725 tip-tilt imaging in Ly α at the UH 2.2m telescope and deep narrow-band IR imaging in [O II] at CFHT reveals a complex and structured system which may be merging to produce the host galaxy of the quasar. For BR2237–0607 the Ly α -emitting galaxies show the strong Ly α -emission and weak continuum expected for objects before substantial chemical enrichment and consequent dust formation and extinction has occurred. These objects appear to be galaxies in their first outburst of star formation.

1 Introduction

It is clear that some form of activity took place prior to $z \sim 5$ to generate the ionization of the intergalactic medium¹ and also that some quasar absorption-line systems have small amounts of metals in place² by $z = 4-5$, indicating that some galaxy formation has initiated at or prior to these redshifts. These early epochs may be our best chance to see galaxies in the first stages prior to the formation of substantial dust and metals, and hence may be the best place for Ly α emission-line searches. While earlier Ly α searches were unsuccessful,^{3,4} improved sensitivities – particularly with the Keck 10m telescope – are now enabling us to detect Ly α emitters at $z \sim 4.5-5$, at least around the known $z > 4.5$ quasars, and there have been a flurry of recent results.^{5,9,6,7} Parallel searches for color-selected objects are also taking place, tuned to these redshift ranges, which are designed to turn up more evolved objects.^{8,9,10}

We discuss here several high-redshift galaxies which have been identified in deep narrow-band Ly α imaging of the fields surrounding the quasars BR1202–

0725 and BR2237–0607, and studied with followup spectroscopy using the LRIS spectrograph on Keck. These objects appear to follow very closely our expectations for early galaxies although, consistent with objects at lower redshifts, they are not particularly luminous.

2 The companions to BR1202–0725

Both BR1202–0725 and BR2237–0607 were identified in the APM quasar survey^{11,12} and show $z > 4$ damped Ly α (DLA) systems in their spectra.¹³ The damped system at $z = 4.383$ [refs. 2, 13] seen in BR1202–0725 is the highest redshift DLA known. Narrow-band imaging⁵ by Hu, McMahon, and Egami at the wavelength of redshifted Ly α for BR1202–0725 identified emission which coincides with the continuum light from a faint galaxy seen in Hubble Space Telescope WFPC2 images 2.6'' NW of the quasar, and pointing radially towards it. This object was also identified as a high-redshift galaxy by Fontana *et al.* using color-break techniques⁹ and by Petitjean *et al.* using integral field spectroscopy.⁶ From high-resolution tip-tilt imaging (FWHM 0.45'') of this system at the UH 2.2m telescope in the Ly α line we have now identified a second faint Ly α emission component coincident with a fainter galaxy 3'' SW of the quasar, also pointing radially towards the quasar in the *HST* image (Fig. 1). In these higher resolution images it is possible to see that the peak of the Ly α emission coincides with the two continuum structures seen in the *HST* images, but that there is also a diffuse component extending out towards the illuminating quasar. The second emission system near the quasar at $z = 4.7$ is confirmed in multi-slit LRIS spectra taken on these objects with the Keck 10m telescope. CIV emission is not detected in these spectra. However, [OII] emission is also seen in these objects in narrow-band IR filter imaging at CFHT made using the UH 1024² IR camera, QUIRC. The morphology of these nearby systems suggests that we may be seeing here the merging of these subsystems and possibly witnessing the formation of the quasar's host galaxy. Near these emission-line structures there is excess emission at millimeter and submillimeter wavelengths^{14,15} and most recently, molecular CO transitions have been detected in close proximity to the quasar.¹⁶

3 Field Galaxies around BR2237–0607 — Early Star Formation?

For BR2237–0607 based on deep narrow-band Ly α imaging followed by spectroscopy using LRIS at Keck, we identify⁷ two Ly α emitting galaxies at $z = 4.55$. In contrast to the BR1202–0725 emission-line objects, these galaxies have separations $> 100''$ (~ 700 kpc) from the quasar, which is unlikely to

Figure 1: The left-hand panel shows a deep *HST* WFPC2 image of the BR1202–0725 field with faint neighboring continuum structures circled ($3''$ diameter). The right-hand panel shows the corresponding Ly α image taken in a 4-hr integration through an $80 \text{ \AA}$ -wide narrow-band filter at the f/31 tip-tilt secondary of the UH 2.2m telescope. Diffuse Ly α emission was detected coincident with both NW and SW companion galaxies to the quasar.

have a significant role in exciting them. Observed equivalent widths for the Ly α line exceed $\sim 1000 \text{ \AA}$, in each case. While one of these objects is quite compact, and has no detectable continuum in a 1-hr *I*-band exposure at Keck, and thus might be an AGN, the second has a diffuse extended structure in both continuum and emission, and is most likely a star-forming galaxy. Line fluxes are $\sim 5 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$. These objects may represent the very earliest stages of galaxy formation prior to the formation of a significant stellar population and the destruction of the Ly α line by dust.

4 Discussion

The objects in the BR2237–0607 field are currently the only galaxies, well separated from quasars, that are known at these high redshifts, and their properties are therefore of considerable interest if only in targeting future searches. The rest frame equivalent widths of the systems are around $> 130 \text{ \AA}$ and $> 240 \text{ \AA}$, which are marginally consistent with stellar excitation for an initial mass function dominated by massive stars.¹⁹ The Ly α may arise from a combination of internal and external ionization. If the observed emission were primarily due to stars, and there were no internal scattering and extinction, then the luminosity of $3 \times 10^{42} h^{-2} \text{ erg s}^{-1}$ ($q_0=0.5$) would correspond to a star formation rate in solar masses ($M_\odot$) per year of $\sim 3 h^{-2} M_\odot \text{ yr}^{-1}$, where we use Kennicutt’s (ref. 21) relation between H α luminosity and star formation rate (SFR) of $\text{SFR} = L(\text{H}\alpha) \times 8.9 \times 10^{-42} \text{ erg s}^{-1} M_\odot \text{ yr}^{-1}$, and assume a ratio of Ly α to H α (8.7) that applies for Case B recombination.²⁰ Given the Hubble time at this redshift of $7 \times 10^8 h^{-1} \text{ yr}$ ($q_0=0.5$) the integrated amount of star formation is small compared to that of a ‘normal’ galaxy with $6 \times 10^{10} h^{-1} M_\odot$ of stars (a so-called L^* galaxy).

Because of the targeted nature of the search it is hard to estimate from the present data whether such objects may be common in the general field or whether they are preferentially found around quasars. Observations, currently in progress, of additional quasars and blank field regions should answer this question.

Acknowledgments

EMH gratefully acknowledges the support of a University Research Council Seed Money grant and grant GO-5975 from STScI. RGM acknowledges the Royal Society.

References

- [1] M.J. Rees, these proceedings.
- [2] L. Lu, *et al.*, ApJ, 457, 1, 1996.
- [3] C.J. Pritchett, PASP, 106, 1052, 1994.
- [4] D. Thompson, S. Djorgovski and J. Trauger, AJ, 110, 963, 1995.
- [5] E.M. Hu, R.G. McMahon and E. Egami, ApJ, 459, L53, 1996.
- [6] P. Petitjean, *et al.*, Nature, 380, 411, 1996.
- [7] E.M. Hu and R.G. McMahon, Nature, 382, 231, 1996.
- [8] E. Egami, PhD thesis, Univ. of Hawaii, 1995.
- [9] A. Fontana, *et al.*, MNRAS, 279, L27, 1996.
- [10] S. D'Odorico *et al.*, these proceedings.
- [11] M.J. Irwin, R.G. McMahon and C. Hazard, in *The Space Distribution of Quasars*, ed. D. Crampton, 117, 1996.
- [12] L.J. Storrie-Lombardi, R.G. McMahon, M.J. Irwin and C. Hazard, ApJ in press, [astro-ph/9604021].
- [13] L.J. Storrie-Lombardi, M.J. Irwin and R.G. McMahon, MNRAS in press, [astro-ph/9608146].
- [14] R.G. McMahon, *et al.*, MNRAS, 267, 9L, 1994.
- [15] K.G. Isaak, *et al.*, MNRAS, 267, L28, 1994.
- [16] A. Omont *et al.*, Nature, 382, 428, 1996.
- [17] L.L. Cowie, A. Songaila, E.M. Hu and J.G. Cohen, AJ, in press.
- [18] L.L. Cowie, E.M. Hu and A. Songaila, Nature, 377, 603, 1995.
- [19] S. Charlot and S.M. Fall, ApJ, 415, 580, 1993.
- [20] M. Brocklehurst, MNRAS, 153, 471, 1971.
- [21] R.C. Kennicutt, Jr., ApJ, 272, 54, 1983.

This figure "tip_tilt.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9609188v1>